

would have been expected. This finding does not prove that LAF can be inherited, but it certainly supports the possibility.

The genetic connection is also supported by work done by Dr. Ramon Brugada and his colleagues at the Baylor College of Medicine and the University of Barcelona[35]. These researchers located three families in Spain in which 21 of 49 family members had lone atrial fibrillation. They mapped their genes and concluded that in these families a mutation in a specific chromosome region (10q22-q24) was the cause of their atrial fibrillation. Dr. Maurits Allesie, MD of the University of Maastricht in the Netherlands makes several very interesting observations concerning these findings[36]:

- If, as in the three Spanish families, lone atrial fibrillation in the general population is also caused by a genetic mutation, then Brugada's findings are of paramount importance.
- The possibility that small molecular defects in DNA can cause changes in the electrophysiologic properties of the atria that, in turn, create a substrate for chronic atrial fibrillation is not unlikely. NOTE: The term chronic in this statement does not mean permanent as opposed to paroxysmal, but rather that LAF is not an acute condition.
- The genes that encode adrenergic receptors are located at the observed mutation site on chromosome 10q. This means that the basis for familial atrial fibrillation could lie in abnormal atrial triggering mechanisms.

Dr. Allesie concludes, "The anatomical and electrophysiologic features of the atria are such that there is only a narrow margin of safety between normal sinus rhythm and chronic atrial fibrillation." Our survey findings of a possible genetic connection and the fascinating discoveries of Dr. Brugada and colleagues together with Dr. Allesie's profound observations certainly provide much food for thought and will hopefully be followed up by additional research.

The finding that some individuals have a genetic abnormality favouring lone atrial fibrillation does not mean that they are predestined to develop LAF, but it could mean that they need to be more vigilant in avoiding common triggers and known risk factors.

## Autonomic Nervous System Imbalances

The idea that the autonomic nervous system (ANS) may be intimately involved in LAF was first put forward in 1982 by Philippe Coumel, MD of the Lariboisiere Hospital in Paris, France. Dr. Coumel proposed that LAF is caused by a particularly sensitive heart tissue, a dysfunction of the autonomic nervous system or both[11]. In the 20 years since Dr. Coumel first outlined his findings, much research has been done in the field and it is now generally accepted that there is a close connection between LAF and an ANS imbalance.

Most, if not all, of the research on the LAF/ANS connection has involved the use of highly sophisticated equipment designed to measure heart rate variability.

Normally the heart beats at a rate of 60 to 80 beats per minute, but the rate can be as low as 50 (in trained athletes) or as high as 100 (during illness or in a highly emotional state) without it being cause for concern. The heart rate is a measure of the number of

contractions made by the left ventricle in a one-minute period and is usually measured on the radial artery near the wrist or on the carotid artery in the neck.

So while the heart rate itself has long been known to be an indicator of the physical and psychological condition of an individual it is only recently that attention has focused on the significance of the beat to beat variability. Research has shown that the interval between one heart beat and the next varies and that this variation is a powerful indicator of the state of the autonomic nervous system.

### **Measurement of Heart Rate Variability**

The variation in the heart beat interval is usually measured via a 5-minute electrocardiogram or a 24-hour Holter monitoring. The original and still commonly used measure for the variation is referred to as SDNN which is the standard deviation of the heart beat intervals, that is, the square root of the variance. Powerful and fast calculation techniques developed in the late 1960s made it possible to extract more sophisticated and valuable information from the heart rate recordings. Most scientific work on heart rate variability now uses power spectral density (PSD) analysis to relate the relatively simple measurement of beat to beat variability to the state of the autonomic nervous system. PSD analysis uses a mathematical technique (fast Fourier transform) to determine how the power (variance in heart beat interval) is distributed across different frequency bands. There is now general agreement that the power in the low frequency band (LF) from 0.04 to 0.15 Hz (cycles/second) is an indication of sympathetic (adrenergic) branch activity and that the power in the high frequency band (HF) from 0.15 to 0.40 Hz is primarily an indication of parasympathetic (vagal) activity. It follows that the ratio of LF/HF is a measure of the balance of the autonomic nervous system with a higher number indicating an excess of adrenergic activity and a lower number indicating an excess of vagal activity. The total power (TP) in the PSD analysis, that is the power obtained by considering the whole frequency spectrum, is another important measure of heart rate variability[37].

### **Significance of Heart Rate Variability**

Not surprisingly, heart rate variability analysis was first used as a measure of heart health. In 1996 researchers involved in the Framingham Heart Study concluded that a lower than normal heart rate variability (HRV) was associated with an increased risk of angina, heart attack, and other cardiac events[38]. Other researchers have found correlations between low HRV and the risk of neuropathy in diabetics and between low HRV and depression[37,39]. There is also a very significant association between low HRV and the risk of dying after suffering a heart attack. It is postulated that decreased HRV correlates with increased sympathetic (adrenergic) or decreased vagal tone, which may predispose to ventricular fibrillation[40]. British researchers report that reduced HRV is a risk factor for atherosclerosis in both diabetic and non-diabetic patients[41]. A group of American and Dutch researchers recently concluded that low HRV not only increases the risk of heart disease especially death from a heart attack, but may indeed precede a number of different diseases. As a matter of fact, they make the profound statement that “sympathetic predominance, as reflected in low HRV and high heart rate, may be indicative of less favorable general health, with HRV being a more sensitive indicator than heart rate”. The researchers also point out that HRV decreases with age, high insulin levels, physical inactivity, smoking, and rapid and shallow breathing[42].

It is clear that low HRV is a predictor of poor health and an increased risk for cardiac death. Does this mean that high HRV is beneficial? Not necessarily. A recent study carried out by Dutch researchers clearly shows that elderly people with a very high HRV have an even higher

risk for sudden cardiac death than do elderly people with a very low HRV. So it would seem that there is an optimum range, the exact limits of which still need to be determined[43].

### **HRV and Atrial Fibrillation**

HRV analysis has been used extensively in the study of atrial fibrillation. LAF episodes can be divided into two groups; those that are preceded by an increase in LF power and a decrease in HF power consistent with an increase in sympathetic (adrenergic) tone, and those that are preceded by a decrease in LF power and an increase in HF power consistent with an increase in parasympathetic (vagal) tone. The changes in HRV are apparent at least 5 minutes before the actual episode[44,45]. It is also clear that most vagal afibbers have no underlying heart disease while adrenergic type afibbers often do. There is also considerable evidence that the number of ectopic (premature) beats increases prior to an episode in patients with lone atrial fibrillation[47].

Atrial fibrillation patients with focal ectopy originating from the pulmonary veins experience a significant increase in HF power and a decrease in LF power during the 20 minutes preceding an episode[47]. This may mean that focal AF is primarily found among vagal afibbers raising the interesting question “Is pulmonary vein ablation likely to be more beneficial for vagal afibbers than for adrenergic ones?”

Other research has established that afibbers who have undergone cardioversion for atrial fibrillation are twice as likely to have another episode if their LF/HF ratio is high[48,49].

### **External Influences on HRV**

It is clear that the autonomic nervous system is dysfunctional in lone atrial fibrillation patients and that the extent of this dysfunction manifests itself in several measures of HRV. HRV is highly influenced by emotional state. Anger and anxiety increases LF power while depression reduces overall HRV[39]. Alcohol consumption increases the LF/HF ratio and the blood level of norepinephrine[50]. Exposure to air pollution (particulates) may reduce HRV[51]. Physical exercise, on the other hand, has been found to increase HRV[52,58]. The ingestion of sugar (glucose) is associated with a significant increase in the LF/HF ratio[54]. Melatonin increases vagal tone (HF power) in men when they are lying down so this may not be the greatest supplement for vagal afibbers[55].

Sleeping position can also affect HRV. Sleeping on the right side increases HF power and minimizes LF power leading to a low LF/HF ratio; norepinephrine secretion is also minimized when lying on the right side[56,57]. This could mean that adrenergic afibbers should preferably sleep on their right side.

The link between exposure to electromagnetic radiation (EMF) and HRV is still controversial. Some studies have found a link while others have not. Researchers at the Midwest Research Institute in Kansas City have found that changes in HRV may occur when exposure to electromagnetic fields is combined with stress[58]. Does this mean that your HRV could change if you sit in front of your computer screen while under stress? I used to get very anxious after using my regular desktop computer for a couple of hours. About a year ago I replaced the monitor with a LCD monitor, and since then, I have experienced no problems at all. The LCD monitor emits no radiation and it does not flicker either. I highly recommend it.

Russian researchers believe that abnormalities in the structure of the spine particularly around the seventh and eighth cervical or the first thoracic segments can result in irritation of the sympathetic nerve bundles passing from the spine into the inner body. This, in turn, could affect the sympathetic side of the autonomic nervous system and cause changes in HRV[59]. This phenomenon may explain why some afibbers, presumably adrenergic, have gotten relief from chiropractic manipulations.

Pharmaceutical drugs can also affect HRV. Propafenone (Rythmol) has been found to decrease HRV and LF/HF ratio and flecainide (Tambocor) decreased HRV in heart attack patients[37]. ACE inhibitors (quinapril) and beta-blockers (metoprolol) both increased HF power and decreased LF power and LF/HF ratio in heart attack patients. It is interesting that the effect of quinapril was most pronounced between 2 and 4 AM, 8 and 11 AM, and 7 and 10 PM. Metoprolol was most effective between 8 AM and 12 noon and between 7 and 10 PM[60].

Obviously there are many external factors that may affect HRV. The question then is what can be done to control HRV and prevent imbalances?

### **Control of HRV**

Breathing oxygen-fortified air may be helpful for adrenergic afibbers, but should not be attempted except with the cooperation of a physician. Research has shown that breathing air containing 31% oxygen markedly increases HF power and reduces LF/HF ratio in congestive heart failure patients[61]. Hyperbaric oxygen therapy is also effective in increasing HF power and reducing LF power and the LF/HF ratio in both healthy subjects and in heart disease and stroke patients[62]. These effects are only temporary, but could perhaps be useful in aborting an impending episode. For longer term effects it seems that the mind itself must be engaged. Transcendental meditation has been found to reduce norepinephrine levels and is probably doing this by reducing LF (adrenergic) power[63].

A team of British, Italian and Polish researchers has discovered that repeating a common mantra or reciting the rosary synchronizes and reinforces inherent cardiovascular rhythms. Their experiment involved 23 healthy adults who were instructed to either recite the rosary (in Latin) or to repeat the mantra “om-mani-padme-om” during a 6-minute test period. To their astonishment, the researchers found that the breathing rate of the participants automatically slowed to 6 cycles per minute from the rate of 14.1 observed during spontaneous breathing. This 10-second cycle (6 inhalations/exhalations per minute) is a very important internal rhythm, which synchronizes respiratory and cardiovascular rhythms as well as sympathetic and vagal outflow. Maintenance of the 10-second cycle also increases calmness and well-being. The researchers speculate that the practice of using mantras to achieve the highly beneficial breathing rate of 6 cycles/minute originated in Tibet and India and was introduced to Europe by the Crusaders where it transmuted into the rosary[64].

The most promising approaches for achieving HRV control may well be Thought Field Therapy and the techniques developed by the Heart Math Institute.

### **Thought Field Therapy**

Thought Field Therapy (TFT) was developed by Dr. Roger Callahan about 20 years ago. It is based on the observation that many psychological and even some physical disorders have their origins in negative thought patterns. The purpose of TFT is to eliminate these thought patterns. The technique involves the stimulation of a sequence of acupuncture points by

tapping on them while focusing on the emotion created by thinking about the problem. The precise sequence of tapping depends on the nature of the problem to be eliminated. Tapping on an acupuncture point delivers mechanical energy to the point which is converted to electrical energy which travels along the appropriate meridians and in doing so collapses the negative thought patterns and thereby elicits healing. TFT is now practiced by thousands of psychologists and other health practitioners around the world and, according to its proponents and healed patients, is highly effective.

Dr. Callahan and other TFT practitioners recently released a series of papers discussing the relationship between TFT and heart rate variability. Their work shows that resolution of a problem automatically brings with it a beneficial change in HRV, that is, excessively low HRVs are increased and excessively high HRVs are decreased[65,66]. Dr. Callahan presents a series of 20 cases where the correlation between a successful TFT session and a beneficial change in HRV is clearly demonstrated[65]. Drs. Pignotti and Steinberg discuss 39 cases involving various disorders such as anxiety, depression, fatigue, attention deficit hyperactivity, obsessions, and anger. All cases were successfully treated with TFT and their resolution was accompanied by a beneficial change in HRV. One of the cases involved a 60-year-old man with atrial fibrillation. Prior to treatment his HRV was excessively high. After treatment it decreased to the normal range and the overall balance of the autonomic nervous system was significantly improved[67].

A very large study of TFT was carried out at Kaiser Behavioral Medicine and Behavioral Health Services in Honolulu. A total of 714 patients were treated for 31 different types of psychological problems. Improvements were observed in all 31 categories and were accompanied by improved HRV and autonomic system balance[63,68]. Relief workers in Kosovo reported a very high rate of success (247 of 249 treatments) using TFT to treat various traumas resulting from the war and ethnic cleansing[69].

Although the evidence for effectiveness of TFT and its strong correlation with HRV improvement is convincing the technique is still rejected by mainstream psychologists[70-72].

TFT has spawned other similar but simpler techniques for problem resolution through tapping on acupuncture points. The "Emotional Freedom Technique" (EFT) was developed by Gary Craig, an electrical engineer. EFT differs from TFT in that it uses a standard tapping sequence on 12 acupuncture points irrespective of the nature of the problem to be solved. EFT is very easy to learn (from a skilled practitioner) and is purported to be highly effective.

### **Heart Math Techniques**

Rollin McCraty and his colleagues at the Heart Math Institute in California have found that emotions are strongly related to HRV. Both anger (negative emotion) and appreciation (positive emotion) increase HRV, but only anger increases the LF/HF ratio as well[73]. McCraty also found that sincere appreciation can significantly increase DHEA levels while decreasing cortisol levels[74]. The Heart Math techniques balance the autonomic nervous system (ANS) and achieve a state of coherence between the brain and the heart. The Freeze-Frame technique deals with immediate disturbances in the ANS while the Heart Lock-In technique is aimed at establishing coherence and balancing the ANS on a long-term basis. Both techniques are very simple.

The idea behind Freeze-Frame is to recognize the stressful feeling, shifting your focus to the heart region, and then recalling a positive feeling or experience to “crowd out” the stressful feeling[74].

The Heart Lock-In technique is meant to be practiced on a daily basis. It is quite similar to meditation and has 5 steps much like the Freeze-Frame technique

The Heart Math techniques are very powerful. A clinical trial carried out at the Kaiser Permanente Clinic found that 60 out of 75 atrial fibrillation patients improved markedly and 14 were able to get off all antiarrhythmic medications after practicing the two Heart Math techniques[75].

The Heart Math Institute has a computer program (Freeze Framer) available that will help evaluate progress in using the techniques. However, the program is not necessary in order to apply the techniques effectively.

It is clear there is a strong association between a dysfunctional autonomic nervous system (ANS) and lone atrial fibrillation. Analysis of heart rate variability (HRV) is an excellent method for determining the degree and direction of the ANS abnormality. HRV, in turn, is affected by many external factors and, it would appear, can possibly be changed by the use of new energy psychology techniques such as Freeze-Frame, Heart-Lock-in, TFT and EFT.

I have tried EFT, but did not find it beneficial for my afib. At least one afibber tried TFT, but again did not find it beneficial.

*Note: Heart Math, Freeze-Frame, and Heart Lock-In are registered trademarks of the Institute of Heart Math. Freeze Framer is a trademark of Quantum Intech.*

## References

1. ACC/AHA/ESC Guidelines for the Management of Patients with Atrial Fibrillation: Executive Summary. *Journal of the American College of Cardiology*, Vol. 38, No. 4, October 2001
2. Kottkamp, M.D. et al. Atrial fibrillation: epidemiology, etiology, and symptoms, Chapter 9. In *Atrial Flutter and Fibrillation: From Basic to Clinical Applications*, edited by N. Saoudi, et al. Armonk, NY, Futura Publishing, 1998
3. Havranek, Edward P. The management of atrial fibrillation: current perspectives. *American Family Physician*, Vol. 50, No. 5, October 1994, pp. 959-68
4. Hurst's *The Heart*, 10<sup>th</sup> edition, NY, McGraw-Hill, 2001, pp. 1029-30
5. Garrett, Michael M. Practical management of atrial fibrillation. *Postgraduate Medicine*, Vol. 87, January 1990, pp. 40-49
6. Falk, Rodney H. Proarrhythmic responses to atrial antiarrhythmic therapy, Chapter 17. In *Atrial Fibrillation: Mechanisms and Management*, edited by R.H. Falk and P.J. Podrid, 2<sup>n</sup> ed., Philadelphia, PA, Lippincott-Raven Publishers, 1997
7. Kerr, Charles R. and Leather, Richard A. Atrial fibrillation in the absence of overt cardiac disease, Chapter 9. In *Atrial Fibrillation: Mechanisms and Management*, edited by R.H. Falk and P.J. Podrid, 2<sup>n</sup> ed., Philadelphia, PA, Lippincott-Raven Publishers, 1997
8. Gilligan, David M., et al. The management of atrial fibrillation. *American Journal of Medicine*, Vol. 101, October 1996, pp. 413-21
9. Baer, Margaret, and Goldschlager, Nora. Atrial fibrillation: an update on new management strategies. *Geriatrics*, Vol. 50, April 1995, pp. 22-29
10. Chugh, Sumeet, S., et al. Epidemiology and natural history of atrial fibrillation: clinical implications. *Journal of the American College of Cardiology*, Vol. 37, February 2001, pp. 371-78

11. Coumel, Philippe. The role of the autonomic nervous system in atrial flutter and fibrillation: clinical findings, Chapter 6. In *Atrial Flutter and Fibrillation: From Basic to Clinical Applications*, edited by N. Saoudi, et al. Armonk, NY, Futura Publishing, 1998
12. Thornton, J.R. Atrial fibrillation in healthy non-alcoholic people after an alcoholic binge. *Lancet*, Vol. 2, November 3, 1984, pp. 1013-15
13. Takauchi, Y., et al. Tyramine-induced endogenous noradrenaline efflux from in situ cardiac sympathetic nerve ending in cats. *Acta Physiologica Scandinavica*, Vol. 168, February 2000, pp. 287-93
14. Jacob, L.H. and Carron, D.B. Atrial fibrillation precipitated by tyramine containing foods. *British Heart Journal*, Vol. 57, February 1987, pp. 205-06
15. Leira, R. and Rodriguez, R. Diet and migraine [English abstract only]. *Rev Neurol*, Vol. 24, No. 129, May 1996, pp. 534-38 [article in Spanish]
16. Stein, Jay H., editor. *Internal Medicine*, 3<sup>rd</sup> edition, Boston, MA, Little, Brown and Company, 1990, pp. 2166-83
17. *Harrison's Principles of Internal Medicine*, 12<sup>th</sup> edition, 1991, McGraw-Hill, NY, pp. 1692-1706
18. Burggraaf, J., et al. Sympathovagal imbalance in hyperthyroidism. *American Journal of Physiol Endocrinol Metab*, Vol. 281, No. 1, July 2001, pp. E190-95
19. Auer, J., et al. Subclinical hyperthyroidism as a risk factor for atrial fibrillation. *American Heart Journal*, Vol. 142, No. 5, November 2001, pp. 838-42
20. Sawin, C.T. Subclinical hyperthyroidism and atrial fibrillation. *Thyroid*, Vol. 12, June 2002, pp. 501-03
21. Petretta, M., et al. Cardiovascular haemodynamics and cardiac autonomic control in patients with subclinical and overt hyperthyroidism. *European Journal of Endocrinology*, Vol. 145, No. 6, December 2001, pp. 691-96
22. Siddiqui, A.S., et al. Covert hypothyroidism with weight loss and atrial fibrillation. *British Journal of Clinical Practice*. Vol. 47, No. 5, Sept-Oct. 1993, p. 268
23. Gerritsen, R.J., et al. Relationship between atrial fibrillation and primary hypothyroidism in the dog. *Vet. Quarterly.*, Vol. 18, No. 2, June 1996, pp. 49-51
24. Cacciatori, V., et al. Power spectral analysis of heart rate in hypothyroidism. *European Journal of Endocrinology*, Vol. 143, No. 3, September 2000, pp. 327-33
25. Lerch, M., et al. Is there a need for treatment in subclinical hypo- and hyperthyroidism? [English abstract only]. *Ther Umsch*, Vol. 56, No. 7, July 1999, pp. 369-73 [article in German]
26. Odeh, Majed, et al. Transient atrial fibrillation precipitated by hypoglycemia. *Annals of Emergency Medicine*, Vol. 19, May 1990, pp. 565-67
27. Yinnon, A.M., et al. Hypoglycemia - a rare cause of atrial fibrillation. *Isr J Med Sci*, Vol. 25, 1989, pp. 346-47
28. Palardy, Jean, et al. Blood glucose measurements during symptomatic episodes in patients with suspected postprandial hypoglycemia. *New England Journal of Medicine*, Vol. 321, November 23, 1989, pp. 1421-25
29. Portis, Sidney A. Life situations, emotions and hyperinsulinism. *Journal of the American Medical Association*, Vol. 142, April 22, 1950, pp. 1281-86
30. D'Alessio, D.A., et al. Activation of the parasympathetic nervous system is necessary for normal meal-induced insulin secretion in rhesus macaques. *J Clin Endocrinol Metab*, Vol. 86, March 2001, pp. 1253-59
31. Stein, Jay H., editor. *Internal Medicine*, 3<sup>rd</sup> edition, Boston, MA, Little, Brown and Company, 1990, p. 855-60
32. *Harrison's Principles of Internal Medicine*, 12<sup>th</sup> edition, 1991, McGraw-Hill, NY, pp. 285-87
33. Kottkamp, Hans, et al. Atrial fibrillation: epidemiology, etiology, and symptoms, Chapter 9. *From Atrial Flutter and Fibrillation*, edited by N. Saoudi, et al. Armonk, NY, Futura Publishing Company, 1998, pp. 135-51
34. Lenders, J.W., et al. Biochemical diagnosis of pheochromocytoma: which test is best? *Journal of the American Medical Association*, Vol. 287, March 20, 2002, pp. 1427-34
35. Brugada, Ramon, et al. Identification of a genetic locus for familial atrial fibrillation. *New England Journal of Medicine*, Vol. 336, March 27, 1997, pp. 905-11
36. Allessie, Maurits A. Is atrial fibrillation sometimes a genetic disease? *New England Journal of Medicine*, Vol. 336, March 27, 1997, pp. 950-52

37. Heart rate variability: standards of measurement, physiological interpretation, and clinical use. *Circulation*, Vol. 93, March 1, 1996, pp. 1043-65
38. Tsuji, Hisako, et al. Impact of reduced heart rate variability on risk of cardiac events. *Circulation*, Vol. 94, December 1, 1996, pp. 2850-55
39. Carney, Robert M., et al. Association of depression with reduced heart rate variability in coronary artery disease. *American Journal of Cardiology*, Vol. 76, September 15, 1995, pp. 562-64
40. Kleiger, Robert E., et al. Decreased heart rate variability and its association with increased mortality after acute myocardial infarction. *American Journal of Cardiology*, Vol. 59, February 1, 1987, pp. 256-62
41. Colhoun, Helen M., et al. The association of heart-rate variability with cardiovascular risk factors and coronary artery calcification. *Diabetes Care*, Vol. 24, June 2001, pp. 1108-14
42. Dekker, Jacqueline M., et al. Low heart rate variability in a 2-minute rhythm strip predicts risk of coronary heart disease and mortality from several causes. *Circulation*, Vol. 102, September 12, 2000, pp. 1239-44
43. de Bruyne, Martine C., et al. Both decreased and increased heart rate variability on the standard 10-second electrocardiogram predict cardiac mortality in the elderly. *American Journal of Epidemiology*, Vol. 150, No. 12, 1999, pp. 1282-88
44. Fioranelli, M., et al. Analysis of heart rate variability five minutes before the onset of paroxysmal atrial fibrillation. *Pacing Clin Electrophysiol*, Vol. 22, May 1999, pp. 743-49
45. Huang, J.L., et al. Changes of autonomic tone before the onset of paroxysmal atrial fibrillation. *International Journal of Cardiology*, Vol. 66, October 30, 1998, pp. 275-83
46. Vikman, Saila, et al. Altered complexity and correlation properties of R-R interval dynamics before the spontaneous onset of paroxysmal atrial fibrillation. *Circulation*, Vol. 100, November 16, 1999, pp. 2079-84
47. Zimmermann, M. and Kalusche, D. Fluctuation in autonomic tone is a major determinant of sustained atrial arrhythmias in patients with focal ectopy originating from the pulmonary veins. *Journal of Cardiovascular Electrophysiology*, Vol. 12, March 2001, pp. 285-93
48. Lombardi, Federico, et al. Heart rate variability and early recurrence of atrial fibrillation after electrical cardioversion. *Journal of the American College of Cardiology*, Vol. 37, January 2001, pp. 157-62
49. Michelucci, A., et al. Altered values of heart rate variability in patients with relapse of atrial fibrillation during the first week after electrical cardioversion: preliminary data. *Italian Heart Journal*, Vol. 2, June 2001, pp. 435-40
50. Maki, T., et al. Effect of ethanol drinking, hangover, and exercise on adrenergic activity and heart rate variability in patients with a history of alcohol-induced atrial fibrillation. *American Journal of Cardiology*, Vol. 82, August 1, 1998, pp. 317-22
51. Magari, Shannon R., et al. Association of heart rate variability with occupational and environmental exposure to particulate air pollution. *Circulation*, Vol. 104, August 28, 2001, pp. 986-91
52. Schuit, Albertine J., et al. Exercise training and heart rate variability in older people. *Medicine & Science in Sports & Exercise*, Vol. 31, June 1999, pp. 816-21
53. Masari, Amano, et al. Exercise training and autonomic nervous system activity in obese individuals. *Medicine & Science in Sports & Exercise*, Vol. 33, August 2001, pp. 1287-91
54. Paolisso, G., et al. Effects of glucose ingestion on cardiac autonomic nervous system in healthy centenarians: differences with aged subjects. *European Journal of Clin Invest*, Vol. 30, April 2000, pp. 277-84
55. Nishiyama, K., et al. Acute effects of melatonin administration on cardiovascular autonomic regulation in healthy men. *American Heart Journal*, Vol. 141, May 2001, p. E9
56. Kuo, Cheng-Deng and Chen, Gau-Yang. Comparison of three recumbent positions on vagal and sympathetic modulation using spectral heart rate variability in patients with coronary artery disease. *American Journal of Cardiology*, Vol. 81, February 15, 1998, pp. 392-96
57. Fujita, M., et al. Effects of posture on sympathetic nervous modulation in patients with chronic heart failure. *Lancet*, Vol. 356, November 25, 2000, pp. 1822-23
58. Graham, C., et al. Cardiac autonomic control mechanisms in power-frequency magnetic fields: a multistudy analysis. *Environ Health Perspect*, Vol. 108, August 2000, pp. 737-42



59. Mironova, T.F. and Mironov, V.A. Clinical analysis of heart rate variability. Paper presented at the First Virtual HRV Congress held in Kiev, Ukraine, November 2001 (<http://www.hrvcongress.org>)
60. Kontopoulos, A.G., et al. Effect of quinapril or metoprolol on circadian sympathetic and parasympathetic modulation after acute myocardial infarction. *American Journal of Cardiology*, Vol. 84, November 15, 1999, pp. 1164-69
61. Bartels, M.N., et al. Oxygen supplementation and cardiac-autonomic modulation in COPD. *Chest*, Vol. 118, September 2000, pp. 691-96
62. Stepanov, Andrey and Stepanova, Svetlana. Heart rate variability in patients during hyperbaric oxygenation. Paper presented at the First Virtual HRV Congress held in Kiev, Ukraine, November 2001
63. Infante, J.R., et al. Catecholamine levels in practitioners of transcendental meditation technique. *Physiol Behav*, Vol. 72, January 2001, pp. 141-46
64. Bernardi, Luciano, et al. Effect of rosary prayer and yoga mantras on autonomic cardiovascular rhythms: comparative study. *British Medical Journal*, Vol. 323, December 22-29, 2001, pp. 1446-49
65. Callahan, Roger J. The impact of Thought Field Therapy on heart rate variability. *Journal of Clinical Psychology*, Vol. 57, October 2001, pp. 1153-70
66. Callahan, Roger J. Raising and lowering of heart rate variability: some clinical findings of Thought Field Therapy. *Journal of Clinical Psychology*, Vol. 57, October 2001, pp. 1175-86
67. Pignotti, Monica and Steinberg, Mark. Heart rate variability as an outcome measure for Thought Field Therapy in clinical practice. *Journal of Clinical Psychology*, Vol. 57, October 2001, pp. 1193-1206
68. Sakai, Caroline, et al. Thought Field Therapy clinical applications: utilization in an HMO in behavioural medicine and behavioural health services. *Journal of Clinical Psychology*, Vol. 57, October 2001, pp. 1215-27
69. Johnson, Carl, et al. Thought Field Therapy: soothing the bad moments of Kosovo. *Journal of Clinical Psychology*, Vol. 57, October 2001, pp. 1237-40
70. McNally, Richard J. Tertullian's motto and Callahan's method. *Journal of Clinical Psychology*, Vol. 57, October 2001, pp. 1171-74
71. Herbert, James D. and Gaudiano, Brandon A. The search of the Holy Grail: heart rate variability and Thought Field Therapy. *Journal of Clinical Psychology*, Vol. 57, October 2001, pp. 1207-14
72. Rosner, Rita. Between search and research: how to find your way around? Review of the article "Thought Field Therapy – soothing the bad moments of Kosovo". *Journal of Clinical Psychology*, Vol. 57, October 2001, pp. 1241-44
73. McCraty, Rollin, et al. The effects of emotions on short-term power spectrum analysis of heart rate variability. *American Journal of Cardiology*, Vol. 76, November 15, 1995, pp. 1089-93
74. Childre, Doc and Martin, Howard. *The Heart Math Solution*. HarperCollins Publishers, NY, 1999
75. Personal communication with Rollin McCraty, November 15, 2001
76. Dayan, Colin M., et al. Whose normal thyroid function is better – yours or mine? *The Lancet*, Vol. 360, August 3, 2002, pp. 353-54
77. Elisaf, M., et al. Hypokalaemia in alcoholic patients. *Drug Alcohol Rev*, Vol. 21, No. 1, March 2002, pp. 73-6
78. Oe, H., et al. A case of severe verapamil overdose. *Japan Circ J*, Vol. 62, Jan. 1998, pp. 72-6
79. Schwinger, R.H. and Erdmann, E. Heart failure and electrolyte disturbances. *Methods Find Exp Clin Pharmacol*, Vol. 14, No. 4, May 1992, pp. 315-25
80. Kruse, J.A., et al. Concentrated potassium chloride infusions in critically ill patients with hypokalemia. *J Clin Pharmacol*, Vol. 34, November 1994, pp. 1077-82